

## Reprint acknowledgements

Figure 2.12: Reprinted with permission from *J. Am. Chem. Soc.*, vol. 91; T. Williams; R. G. Pitcher; P. Bommer; J. Gutzwiller; M. Uskokovic, “*Diastereomeric Solute-Solute Interactions of Enantiomers in Achiral Solvents. Nonequivalence of the Nuclear Magnetic Resonance Spectra of Racemic and Optically Active Dihydroquinine*”, pages 1871-1872 (1969). Copyright 1969, American Chemical Society.

Figure 4.6b: Reprinted with permission from *J. Am. Chem. Soc.*, vol. 95; H. B. Bürgi; J. D. Dunitz; E. Scheffter, “*Geometrical Reaction Coordinates. II. Nucleophilic Addition to a Carbonyl Group*”, pages 5065-5067 (1973). Copyright 1973 American Chemical Society.

Figure 4.7: Reprinted from *Tetrahedron*, vol. 30; H. B. Bürgi; D. Dunitz; J. M. Lehn; G. Wipff, “*Stereochemistry of Reaction Paths at Carbonyl Centres*”, pages 1563-1572 (1974), with kind permission from Elsevier Science Ltd., The Boulevard, Langford Lane, Kidlington OX5 1GB, UK.

Figure 4.9a: Reprinted with permission from *J. Am. Chem. Soc.*, vol. 109; E. P. Lodge; C. H. Heathcock, “*On the Origin of Diastereofacial Selectivity in Additions to Chiral Aldehydes and Ketones: Trajectory Analysis*”, 2819-2820 (1987). Copyright 1987 American Chemical Society.

Figure 6.14: Reprinted with kind permission of VCH Verlagsgesellschaft and Professor G. Helmchen from “*Concerning the Mechanism of the Asymmetric Diels-Alder Reaction: First Crystal Structure Analysis of a Lewis Acid Complex of a Chiral Dienophile*”, by T. Poll; J. O. Metter; G. Helmchen, *Angew. Chem. Int. Ed. Engl.* **1985**, 24, 112-114.

Figure 7.12 (lower part) and Figure 7.13c: Reprinted from *Tetrahedron*, vol. 30; K. N. Houk; N. G. Rondan; Y.-D. Wu; J. T. Metz; M. N. Paddon-Row, “*Theoretical Studies of Stereoselective Hydroborations*”, pages 2257-2274 (1984), with kind permission from Elsevier Science Ltd., The Boulevard, Langford Lane, Kidlington OX5 1GB, UK.